

Important Concepts . . .

# Preview Review



**Mathematics Grade 8**

**W2 - Lesson 2: Solving Linear Equations**

## Important Concepts of Grade 8 Mathematics

W1 - Lesson 1 .....	Perfect Squares and Square Roots
W1 - Lesson 2 .....	Working with Ratios and Rates
W1 - Lesson 3 .....	Multiplying and Dividing Fractions
W1 - Lesson 4 .....	Multiplying and Dividing Integers
W1 - Lesson 5 .....	Working with Percents
W1 - Review	
W1 - Quiz	
W2 - Lesson 1 .....	Modelling and Solving Linear Equations Using Algebra Tiles
W2 - Lesson 2 .....	Solving Linear Equations
W2 - Lesson 3 .....	Graphing and Analyzing Linear Relations
W2 - Lesson 4 .....	Critiquing the Representation of Data
W2 - Lesson 5 .....	Probability of Independent Events
W2 - Review	
W2 - Quiz	
W3 - Lesson 1 .....	Pythagorean Theorem
W3 - Lesson 2 .....	Calculating Surface Area
W3 - Lesson 3 .....	Calculating Volume
W3 - Lesson 4 .....	Drawing 3-D Objects
W3 - Lesson 5 .....	Congruence of Polygons
W3 - Review	
W3 - Quiz	

## Materials Required

Protractor  
Ruler  
Calculator

**No Textbook Required**

**This is a stand-alone course.**

### Mathematics Grade 8

Version 6

Preview/Review W2 - Lesson 2

ISBN 1-891894-00-6

**Publisher:** Alberta Distance Learning Centre

**Written by:** Monica dHamrait

**Reviewed by:** Patty Rogerson

**Project Coordinator:** Donna Silgard

**Preview/Review Publishing Coordinating Team:**

**Heather Martel and Nicole Mckeand**



Alberta Distance Learning Centre has an Internet site that you may find useful. The address is as follows: <http://www.adlc.ca>

The use of the Internet is optional. Exploring the electronic information superhighway can be educational and entertaining. However, be aware that these computer networks are not censored. Students may unintentionally or purposely find articles on the Internet that may be offensive or inappropriate. As well, the sources of information are not always cited and the content may not be accurate. Therefore, students may wish to confirm facts with a second source.

### ALL RIGHTS RESERVED


Copyright © 2010, by Alberta Distance Learning Centre, 4601-63 Avenue, Barrhead, Alberta, Canada, T7N 1P4. Additional copies may be obtained from Alberta Distance Learning Centre.

No part of this courseware may be reproduced or transmitted in any form, electronic or mechanical, including photocopying (unless otherwise indicated), recording, or any information storage and retrieval system, without the written permission of Alberta Distance Learning Centre.

Every effort has been made both to provide proper acknowledgement of the original source and to comply with copyright law. If cases are identified where this effort has been unsuccessful, please notify Alberta Distance Learning Centre so that appropriate corrective action can be taken.

**IT IS STRICTLY PROHIBITED TO COPY ANY PART OF THESE MATERIALS UNDER THE TERMS OF A LICENCE FROM A COLLECTIVE OR A LICENSING BODY.**

# Preview/Review Concepts for Grade Eight Mathematics



*W2 - Lesson 2:*

*Solving Linear Equations*

# OBJECTIVES

By the end of this lesson, you will be able to:

- Solve a one-step and two-step linear equation
- Verify the solution to a given linear equation
- Apply the distributive property to solve a given linear equation
- Solve a given problem using a one-step or two-step linear equation

## GLOSSARY

**Coefficient** – the number located directly in front of the variable

**Constant** – a whole number that is either added to or subtracted from the variable

**Distributive Property** – a mathematical property that states that a product can be written as a sum or difference of two products,  $a(b + c) = ab + ac$  or  $a(b - c) = ab - ac$

**Inverse Operations** – an operation that reverses the effect of another operation. The inverse of addition is subtraction, the inverse of subtraction is addition, the inverse of multiplication is division, and the inverse of division is multiplication.

**Linear Equation** – equations that form a straight non-vertical and non-horizontal line when graphed

**One-step Equations** – equations that can be solved in one-step

**Two-step Equations** – equations that involve two steps in order to solve them

**Variable** – a symbol (usually a letter) that represents an unknown value or values

## W2 - Lesson 2: Solving Linear Equations

### Materials required:


- Paper, and Pencil

### Part 1: Solving One-Step Linear Equations

When solving one-step equations, the goal is to isolate the variable. In order to do this, you must apply the inverse operation to both sides of the equation. Verify your solution by substituting the result back into the equation and evaluating. Ensure that the left side of the equation equals the right side of the equation.

#### Example 1

Solve for  $n$  in the following equation  $3n = -12$ .  
Apply the inverse operation to both sides of the equation to remove the coefficient.

$$\begin{array}{l}
 3n = -12 \\
 \cancel{3}n = \frac{-12}{\cancel{3}} \\
 n = -4
 \end{array}$$


The inverse of multiplication is division.

Divide both sides of the equation by 3.


Verify the solution.

$$\begin{array}{l}
 3n = -12 \\
 3(-4) = -12 \\
 -12 = -12 \\
 \text{LS} = \text{RS}
 \end{array}$$

**Example 2**

Solve for  $r$  in the following equation  $\frac{r}{5} = 8$ .

Apply the inverse operation to both sides of the equation to isolate the variable.



$\frac{r}{5} = 8$

$\cancel{5} \left( \frac{r}{\cancel{5}} \right) = (8)5$

$r = 40$

The inverse of division is multiplication.

Multiply both sides of the equation by 5.

Verify the solution.

$$\begin{aligned}\frac{r}{5} &= 8 \\ \frac{40}{5} &= 8 \\ 8 &= 8 \\ \text{LS} &= \text{RS}\end{aligned}$$

**Practice Questions**

1.  $-13p = 39$

2.  $\frac{b}{8} = 4$

3.  $-10y = -120$

4.  $\frac{s}{-10} = 15$

## Part 2: Solving Two-Step Linear Equations

Solving two-step equations involves a process similar to that of solving one-step linear equations. The main goal is to isolate the variable.

The first inverse operation will involve adding or subtracting the constant from the term that contains the variable.

The second inverse operation will involve multiplication or division to remove the numerical coefficient from the variable.

Remember to apply the inverse operation to both sides of the equation.

Verify your solution by substituting the result back into the equation and evaluating. Ensure that the left side of the equation equals the right side of the equation.



**Example 1**

Solve for  $n$  in the following equation  $5n - 1 = 9$ .

Apply the first inverse operation to remove the constant.

$$5n - 1 = 9$$



$$5n - 1 + 1 = 9 + 1$$

$$5n = 10$$



Apply the second inverse operation to remove the numerical coefficient.

$$5n = 10$$

$$\frac{5n}{5} = \frac{10}{5}$$

$$n = 2$$



verify the solution.

$$5n - 1 = 9$$

$$5(2) - 1 = 9$$

$$10 - 1 = 9$$

$$9 = 9$$

$$LS = RS$$

**Example 2**

Solve for  $b$  in the following equation  $\frac{b}{-2} + 9 = 13$ .

Apply the first inverse operation to remove the constant.

$$\frac{b}{-2} + 9 = 13$$

The inverse of addition is subtraction.

$$\frac{b}{-2} + 9 - 9 = 13 - 9$$

$$\frac{b}{-2} = 4$$

Subtract 9 from both sides of the equation.

Apply the second inverse operation to remove the numerical coefficient.

$$\frac{b}{-2} = 4$$

The inverse of division is multiplication.

$$(-2)\left(\frac{b}{-2}\right) = (4)(-2)$$

$$b = -8$$

Multiply both sides of the equation by -2.

Verify the solution.

$$\frac{b}{-2} + 9 = 13$$

$$\frac{-8}{-2} + 9 = 13$$

$$4 + 9 = 13$$

$$13 = 13$$

$$LS = RS$$

**Practice Questions**

1.  $\frac{r}{4} - 8 = 2$

2.  $-2b + 4 = -20$

3.  $\frac{z}{4} + 10 = -13$

4.  $12c - 8 = -44$

### Part 3: Applying the Distributive Property

The distributive property states that a product can be written as a sum or difference of two products,  $a(b + c) = ab + ac$  or  $a(b - c) = ab - ac$ . Multiply each term inside the brackets by the term located outside the brackets.

#### Example 1

Simplify the algebraic expression  $3(x + 10)$ .  
Multiply each term inside the brackets by the 3.


$$\begin{aligned} 3(x + 10) &= 3(x) + 3(10) \\ &= 3x + 30 \end{aligned}$$

Sometimes a linear equation will need to be simplified before it can be solved. Apply the distributive property before applying the inverse operations when solving two-step linear equations.

#### Example 2

Solve the linear equation  $4(x - 8) = -24$ .  
Apply the distributive property to remove the brackets.


$$\begin{aligned} 4(x - 8) &= -24 \\ 4(x) - 4(-8) &= -24 \\ 4x + 32 &= -24 \end{aligned}$$



Multiply each term in the brackets by 4.

Apply inverse operations to remove the constant.


$$\begin{aligned} 4x + 32 - 32 &= -24 - 32 \\ 4x &= -56 \end{aligned}$$



Subtract both sides of the equation by 32.

Apply inverse operations to remove the numerical coefficient.

$$\begin{aligned} \frac{4x}{4} &= \frac{-56}{4} \\ x &= -14 \end{aligned}$$



Divide both sides of the equation by 4.

**Practice Questions**

1.  $7(x + 12) = 56$

2.  $-3(x - 10) = 48$

## Part 4: Problem Solving with Linear Equations

When solving problems translate the situation represented in the problem into a linear equation. Then solve for the unknown.

### Example 1

Laura bought 5 hockey sticks for \$65.00. Each stick was on sale for \$8.00 off. How many sticks did Laura purchase in total?

**Step 1:** Translate the situation represented in the problem into a linear equation.

Let  $x$  represent the original price of the hockey sticks.

Each hockey stick was on sale for \$8.00 off and Laura bought 5 of them. This can be expressed as  $5(x - 8)$ .

Laura spent \$65.00 in total. The equation becomes  $5(x - 8) = 65$ .

**Step 2:** Solve for the unknown in the linear equation.

Apply the distributive property.

$$5(x - 8) = 65$$

$$5(x) - 5(8) = 65$$

$$5x - 40 = 65$$

Apply inverse operations to isolate the variable.

$$5x - 40 = 65$$

$$5x - 40 + 40 = 65 + 40$$

$$5x = 105$$

$$\frac{5x}{5} = \frac{105}{5}$$

$$x = 21$$

The hockey sticks were originally priced at \$21.00.

**Example 2**

Rena is 6 years older than twice Aaron's age. How old is Aaron if Rena is 22 years old?

**Step 1:** Translate the situation represented in the problem into a linear equation.

Let  $a$  represent Aaron's age.

Rena is 6 years older than twice Aaron's age can be expressed as  $2a + 6$ .

Rena is 22 years old. The equation becomes  $2a + 6 = 22$ .

**Step 2:** Solve for the unknown in the linear equation.

Apply inverse operations to isolate the variable.

$$2a + 6 = 22$$

$$2a + 6 - 6 = 22 - 6$$

$$2a = 16$$

$$\frac{2a}{2} = \frac{16}{2}$$

$$a = 8$$

Aaron is 8 years old.

**Practice Questions**

1. On a fishing trip, Bethany caught two more fish than three times the number of fish Alex caught. If Bethany caught eight fish, how many did Alex catch?

2. Heather is lifting weights in the exercise room. She puts on 6 equal sized weights on her bar. The total amount Heather is lifting is 84 kg. The clamps that hold each weight in place have a mass of 1 kg, how much does each weight weigh?

**Lesson 7: Assignment**

Solve the following linear equations.

1.  $2a = 20$

2.  $\frac{z}{5} = -16$

3.  $-15x = 45$

4.  $-9t = -27$

5.  $\frac{a}{2} + 1 = -4$

6.  $2x - 7 = 9$

7.  $18 = 6 - 3a$

8.  $\frac{b}{12} - 4 = 1$

9.  $10 + 7r = 31$

10.  $-6 + \frac{a}{3} = 5$

11.  $3(x - 7) = -27$

12.  $-5(v + 9) = -60$

13.  $-4(x - 2) = 36$

14.  $-8(m + 7) = 48$

15. Dale is three times as old as Salima. If Dale is 36, how old is Salima?

16. Maple has \$5.00 less than one third of the money that Chan has. If Maple has \$25.00, how much money does Chan have?
17. Vancouver averages 9 less than 4 times the number of rainy days (with 5 mm or more) than Edmonton does. If Vancouver gets 123 days of rain, how many rainy days does Edmonton get?

18. Jagar had three baseball cards. He bought six more packages. He now has 51 cards. How many cards were in each package if each package contained the same number?
19. In Canada, we measure temperature in degrees Celsius ( $^{\circ}\text{C}$ ). In the United States, temperature is measured in degrees Fahrenheit ( $^{\circ}\text{F}$ ). The temperature measured in Fahrenheit is 32 more than 1.8 times the temperature measured in Celsius. If the temperature in Seattle is  $77^{\circ}\text{F}$ , what would it be expressed as in degrees Celsius?

20. On Saturday, Damaris went to the zoo with three of her friends. The zoo reduces its admission prices on Saturdays by \$4.00. In total, the girls spent \$32.00. What would be the cost of admission had the girls gone to the zoo on Friday?





