

Important Concepts . . .

Preview Review



Mathematics Grade 9 TEACHER KEY
W2 - Review

Important Concepts of Grade 9 Mathematics

W1 - Lesson 1	Powers
W1 - Lesson 2	Exponents
W1 - Lesson 3	Rational Numbers
W1 - Lesson 4	Order of Operations
W1 - Lesson 5	Square Roots of Rational Numbers
W1 - Review	
W1 - Quiz	
W2 - Lesson 6	Graphing Linear Relations
W2 - Lesson 7	Solving Linear Relations
W2 - Lesson 8	Linear Inequalities
W2 - Lesson 9	Polynomials
W2 - Lesson 10	Surface Area of 3D Objects
W2 - Review	
W2 - Quiz	
W3 - Lesson 11	Properties of Circles
W3 - Lesson 12	Polygons and Scale Diagrams
W3 - Lesson 13	Rotational Symmetry
W3 - Lesson 14	Representing Data
W3 - Lesson 15	Probability
W3 - Review	
W3 - Quiz	

Materials Required

Pencil
Paper
Calculator
Algebra Tiles

No Textbook Required

This is a stand-alone course.

Mathematics Grade 9

Version 6

Preview/Review W2 - Review

ISBN: 978-1-927090-00-8

Publisher: Alberta Distance Learning Centre

Written by: Lenee Fyfe

Reviewed by: Danielle Winter

Project Coordinator: Danielle Winter

Preview/Review Publishing Coordinating Team: Julie Reschke



Alberta Distance Learning Centre has an Internet site that you may find useful. The address is as follows: <http://www.adlc.ca>

The use of the Internet is optional. Exploring the electronic information superhighway can be educational and entertaining. However, be aware that these computer networks are not censored. Students may unintentionally or purposely find articles on the Internet that may be offensive or inappropriate. As well, the sources of information are not always cited and the content may not be accurate. Therefore, students may wish to confirm facts with a second source.

ALL RIGHTS RESERVED

Copyright © 2011, by Alberta Distance Learning Centre, 4601-63 Avenue, Barrhead, Alberta, Canada, T7N 1P4. Additional copies may be obtained from Alberta Distance Learning Centre.

No part of this courseware may be reproduced or transmitted in any form, electronic or mechanical, including photocopying (unless otherwise indicated), recording, or any information storage and retrieval system, without the written permission of Alberta Distance Learning Centre.

Every effort has been made both to provide proper acknowledgement of the original source and to comply with copyright law. If cases are identified where this effort has been unsuccessful, please notify Alberta Distance Learning Centre so that appropriate corrective action can be taken.

IT IS STRICTLY PROHIBITED TO COPY ANY PART OF THESE MATERIALS UNDER THE TERMS OF A LICENCE FROM A COLLECTIVE OR A LICENSING BODY.

Preview/Review Concepts for Grade Nine Mathematics

Teacher Key



W2 - Review

W2 - Review

Materials required:

- Paper, Pencil, Calculator, Algebra Tiles

Part 1: Graphing Linear Relations

Many patterns, both pictorial and written can be represented using a table of values. This table of values can also be expressed as a linear equation.

Example 1

x	y
-1	-1
0	2
1	5
2	8
3	11

The equation that represents this table of values is $y = 3x + 2$

Linear relations can be checked by substituting in the value that was solved.

Example 2

Using the above example,

$$y = 3x + 2$$

Choose an “ x ” value from the table of values, for example “3”.

$$y = 3(3) + 2$$

Substitute the value of $x = 3$.

$$y = 9 + 2$$


Check back to the table of values to see if when $x = 3$, does $y = 11$?

Yes. The solution is verified.

Example 3

Linear relations can also be graphed to find unknown values. Given the linear relation $y = 3x + 2$, substitute in values for x to determine the values for y . Fill in the table of values. Then, plot the points on a graph.

$y = 3x + 2$	
x	y
-1	-1
0	2
1	5
2	8
3	11



Extending the line can also determine unknown values. For example, the line would extend to point $(-2, -4)$. This indicates that if $x = -2$, then y would equal -4 .

Part 2: Solving Linear Relations

An equation is a number sentence containing a variable and an = sign .

Some examples of equations are:

$$ax = b \quad \frac{x}{a} = b \quad \frac{a}{x} = b$$

When solving these types of equations, only one step is required.

To verify an answer, simply substitute the solution in the equation, then check to make sure the right side = the left side.

Example 1

$$\begin{aligned}\frac{x}{3} &= 27 \\ \frac{x}{3}(3) &= 27(3) \\ x &= 81\end{aligned}$$

Verification

$$\begin{aligned}\frac{x}{3} &= 27 \\ \frac{(81)}{3} &= 27 \\ 27 &= 27 \\ \text{Left side} &= \text{Right side}\end{aligned}$$

Example 2

There are also equations that require two steps to solve for an unknown variable. These equations can be in the form:

$$ax + b = c \text{ or } \frac{x}{a} + b = c$$

To solve these linear equations isolate the variables. Two-step equations will require two processes to solve for the variable. Using inverse operations will solve for a variable.

- The inverse of addition is subtraction.
- The inverse of division is multiplication.

Solve

$$\begin{aligned} 2x + 2 &= 10 \\ 2x + 2 - 2 &= 10 - 2 \\ 2x &= 8 \\ \frac{2x}{2} &= \frac{8}{2} \\ x &= 4 \end{aligned}$$

To isolate the variable in a two step equation, use the reverse order of operations. Add or subtract first, then multiply or divide.

Verification

$$\begin{aligned} 2x + 2 &= 10 \\ 2(4) + 2 &= 10 \\ 10 &= 10 \end{aligned}$$

Right side = Left side

Example 3

To isolate the variable in equations of the form $a(x + b) = c$, consider the following steps.

$$\begin{aligned}4(f + 5) &= 36 \\4f + 20 &= 36 \\4f + 20 - 20 &= 36 - 20 \\4f &= 16 \\f &= 4\end{aligned}$$

Verification

$$\begin{aligned}4(f + 5) &= 36 \\4(4 + 5) &= 36 \\4(9) &= 36 \\36 &= 36 \\ \text{Right side} &= \text{Left side}\end{aligned}$$

Example 4

The equations shown in this format contain variables on both sides of the equations. To solve for the variable, it must be isolated:

$$\begin{aligned}10d &= 25(d - 30) \\10d &= 25(d - 30) && \text{Use the distributive property.} \\10d &= 25d - 750 \\10d - 25d &= 25d - 25d - 750 \\-15d &= -750 \\ \frac{-15d}{-15} &= \frac{-750}{-15} \\d &= 50\end{aligned}$$

Verification

$$\begin{aligned}10d &= 25(d - 30) \\10(50) &= 25(50 - 30) \\500 &= 25(20) \\500 &= 500 \\ \text{Left side} &= \text{Right side}\end{aligned}$$

Example 5

The equations shown in this format contain fractions. To solve for the variable, first remove all fractions by multiplying each term by the LCD (lowest common denominator).

$$4n - \frac{2}{10} = \frac{3}{5}$$

$$(10)4n - (10)\frac{2}{10} = (10)\frac{3}{5}$$

$$40n - 2 = 6$$

$$40n = 8$$

$$n = 5$$

The lowest common denominator of 10 and 5 is 10. So, multiply each term by 10.

Practice Questions

1. Solve for each of the following. Verify your answers for b and d.

a. $2v = -\frac{5}{6}$

$$(2v)(6) = -5$$

$$12v = -5$$

$$v = -\frac{5}{12}$$

b. $\frac{3}{5} = \frac{a}{4}$

$$\left(\frac{3}{5}\right)(4) = \frac{a}{4}(4)$$

$$\frac{12}{5} = a$$

$$a = 2\frac{2}{5}$$

c. $3x = 21$

$$\frac{3x}{3} = \frac{21}{3}$$

$$x = 7$$

Verification

$$3x = 21$$

$$3(7) = 21$$

$$21 = 21$$

Left side = Right side

b. $\frac{14}{x} = 7$

$$14 = 7x$$

$$\frac{14}{7} = x$$

$$2 = x$$

Verification

$$\frac{14}{x} = 7$$

$$\frac{14}{2} = 7$$

$$7 = 7$$

Left side = Right side

2. Solve for each of the following.

a. $2.05 = 0.9x - 6.5$

$$2.05 + 6.5 = 0.9x - 6.5 + 6.5$$

$$8.55 = 0.9x$$

$$x = 9.5$$

b. $12.4g + 34.3 = 9.5 - 3.1g$

$$12.4g + 3.1g + 34.3 = 9.5 - 3.1g + 3.1g$$

$$15.5g + 34.3 - 34.3 = 9.5 - 34.3$$

$$15.5g = -24.8$$

$$g = -1.6$$

c. $-\frac{4}{5}(q+1) = -3$

$$-\frac{4}{5}q - \frac{4}{5} = -3$$

$$(5) - \frac{4}{5}q + (5) - \frac{4}{5} = (5) - 3$$

$$-4q - 4 = -15$$

$$-4q - 4 + 4 = -15 + 4$$

$$-4q = -11$$

$$q = \frac{11}{4} = 2\frac{2}{3}$$

d. $10.1 = 0.9 + \frac{h}{-2}$

$$(2)10.1 = (2)0.9 + (2)\frac{h}{-2}$$

$$20.2 = 1.8 - h$$

$$20.2 - 1.8 = 1.8 - 1.8 - h$$

$$18.4 = -h$$

$$h = -18.4$$

3. Solve the following two step equations.

a. $\frac{3}{4} - \frac{r}{3} = \frac{7}{8}$

$$\begin{aligned} (24)\frac{3}{4} - (24)\frac{r}{3} &= (24)\frac{7}{8} \\ 18 - 8r &= 21 \\ -8r &= 3 \\ r &= -\frac{3}{8} \end{aligned}$$

b. $2z = 4(2z - 3)$

$$\begin{aligned} 2z &= 8z - 12 \\ -6z &= -12 \\ z &= 2 \end{aligned}$$

c. $\frac{1}{2} + \frac{b}{6} = \frac{5}{12}$

$$\begin{aligned} (12)\frac{1}{2} + (12)\frac{b}{6} &= (12)\frac{5}{12} \\ 6 + 2b &= 5 \\ 2b &= -1 \\ b &= -\frac{1}{2} \end{aligned}$$

d. $4(y - 2) = 3(y + 8)$

$$\begin{aligned} 4y - 8 &= 3y + 24 \\ 4y - 3y - 8 &= 3y - 3y + 24 \\ y - 8 &= 24 \\ y - 8 + 8 &= 24 + 8 \\ y &= 32 \end{aligned}$$

e. $0.25m = 0.75(m + 40)$

$$\begin{aligned} 0.25m &= 0.75m + 30 \\ 0.25m - 0.75m &= 0.75m - 0.75m + 30 \\ -0.5m &= 30 \\ \frac{-0.5}{-0.5m} &= \frac{30}{-0.5m} \\ m &= -60 \end{aligned}$$

f. $5.60p + 18.75 = 46.26 - 2.26p$

$$\begin{aligned} 5.60p + 18.75 - 18.75 &= 46.26 - 18.75 - 2.26p \\ 5.60p &= 27.51 - 2.26p \\ 5.60p + 2.26p &= 27.51 + 2.26p - 2.26p \\ 7.86p &= 27.51 \\ p &= 3.5 \end{aligned}$$

Part 3: Inequalities

An inequality compares a linear expression that may or may not be equal. The symbols used for inequality are:

$>$ means 'greater than'

$<$ means 'less than'


\geq means 'greater than or equal to'

\leq means 'less than or equal to'


For example, $x \leq -5$ means that the unknown value, x , is less than or equal to -5 .

Example 1

An inequality can be represented graphically.



The above line segment beginning at -5 is not coloured in completely; this shows $>$.



The above line segment beginning at 0 is completely coloured in. This represents \leq .

Example 2

Solving an inequality is very similar to solving an equation. The purpose is to isolate the variable to solve for a solution.

$$x - 5 \leq 10$$

$$x - 5 + 5 \leq 10 + 5$$

$$x \leq 15$$

When each side of the inequality is multiplied or divided by a negative number, the inequality sign must be reversed in order for it to be true.

$$-8x \geq 24$$

$$\frac{-8x}{-8} \leq \frac{24}{-8}$$

$$x \leq -3$$

To solve multi-step inequalities, work step by step to isolate the variable, remembering to reverse the inequality sign only when multiplying or dividing by a negative number.

$$5x - 19 < 36$$

$$5x - 19 + 19 < 36 + 19$$


$$5x < 45$$

$$\frac{5x}{5} < \frac{45}{5}$$


$$x < 9$$

Practice Questions

1. Write the inequality that represents the following number lines.



_____ $x > 50$ _____



_____ $x \geq 50$ _____



_____ $x < 50$ _____

2. Solve the following inequalities. Graph the solution for b and c.

a. $x - 7 > 22$


$$x - 7 + 7 > 22 + 7$$

$$x > 29$$

b. $4 < x + 11$

$$4 - 11 < x + 11 - 11$$

$$-7 < x$$




c. $7x < 2x + 30$

$$7x - 2x < 2x - 2x + 30$$

$$5x < 30$$

$$x < 6$$



d. $-12x + 10 > 19 - 4x$

$$-12x + 10 - 10 > 19 - 10 - 4x$$

$$-12x > 9 - 4x$$

$$-12x + 4x > 9 - 4x + 4x$$

$$-8x > 9$$

$$\frac{-8x}{-8} < \frac{9}{-8}$$

$$x < -\frac{9}{8} = -1\frac{1}{8}$$

Part 4: Polynomials

A polynomial is an algebraic expression that is made up of terms containing variables, coefficients, exponents and constants. These terms are connected together by addition and subtraction operations.

Polynomials can be named by the number of terms in the expression.

- Monomial – has one term.
- Binomial – has two terms.
- Trinomial – has three terms.

$7x^2 + 5x - 8$ This expression has three terms so it is a trinomial. $7x^2$ is the term with the highest degree; it has a degree is 2. Therefore, the degree of this polynomial is 2.

Example 1

To add and subtract polynomials, it is important to collect like terms. Like terms differ only by their numerical coefficients.

Examples of like terms are:

$$-9x \text{ and } 14x$$

$$16z^2 \text{ and } -4z^2$$

$$8ab \text{ and } -ab$$

Addition and subtraction equations can be solved algebraically or by using algebra tiles. To solve algebraically:

$$(3x + 2) + (2x - 3)$$

$$= 3x + 2 + 2x - 3$$

$$= 5x + 2 - 3$$

$$= 5x - 1$$

Remove the brackets.

Collect like terms with common variables.

Collect the numerical terms (constants).

Example 2

When finding the difference in polynomials, determine the opposite polynomials. Use the opposite strategy to subtract polynomials.

$$(4x - 3) - (3x + 2)$$

$$(4x - 3) + (-3x - 2) \quad \text{Add the opposite.}$$

$$4x - 3 - 3x - 2 \quad \text{Remove brackets.}$$

$$x - 3 - 2 \quad \text{Collect like terms with common variables.}$$

$$x = -5 \quad \text{Collect the numerical terms (constants).}$$

Example 3

To multiply polynomials, multiply the numerical coefficients and use the exponent rules to multiply the variables.

$$(4x)(5x)$$

$$\begin{array}{c} \curvearrowright \\ (4x)(5x) \end{array} \quad \text{Multiply the coefficients.}$$

$$\begin{array}{c} \curvearrowright \\ (4x)(5x) \end{array} \quad \text{Multiply the variables.}$$

$$= 20x^2$$

Example 4

To divide polynomials algebraically, divide the numerical coefficients and use the exponent rules to divide the variables.

$$\frac{12xy}{4x}$$

$$\begin{array}{c} \curvearrowleft \\ \frac{12xy}{4x} \end{array} \quad \text{Divide the numerical coefficients.}$$

$$\begin{array}{c} \frac{12xy}{4x} \\ \curvearrowleft \end{array} \quad \text{Apply exponent laws to variables.}$$

$$= 3y$$


Example 5

To multiply polynomials algebraically, expand the expression using the distributive property.


$$\begin{aligned}
 &(4x)(-2x + 5) \\
 &\overset{\curvearrowright}{(4x)(-2x + 5)} \quad \text{First, multiply } (4x) \text{ and } (-2x) = -8x^2. \\
 &\overset{\curvearrowright}{(4x)(-2x + 5)} \quad \text{Then multiply } (4x) \text{ and } (5) = 20x. \\
 &= -8x^2 + 20x
 \end{aligned}$$

Example 6

Polynomials can be modelled with algebra tiles.



To model a polynomial like $2x^2 - 5x + 3$, simply select the corresponding tiles:



Practice Questions

1. Use the polynomial $10x^2 - 2x - 4$ to answer the following questions.

a. What type of polynomial is this? **trinomial**

b. What is the degree? **second**

c. List the variable(s). **x**

d. List the exponent(s). **2**

e. List the coefficient(s). **10, -2**

f. List the constant(s). **-4**

2. Solve the following by adding or subtracting the polynomials.

a. $(3x^2 - 8x + 6) + (9x^2 + 4x - 1)$

$$= 12x^2 - 4x + 5$$

b. $(8 - 6v) + (3 - 8v)$

$$= 11 - 14v$$

c. $(9m - 4) - (3 - 8m)$

$$= 9m - 4 + -3 + 8m$$

$$= 17m - 7$$

d. $(2xy - 4x + y) - (7xy - 3x - y)$

$$= 2xy - 4x + y + -7xy + 3x + y$$

$$= -5xy - x + 2y$$

3. Find the product.

a. $(-32x)(-2y)$
 $= 64xy$

b. $(5a)(-4a - 2)$
 $= -20a^2 - 10a$

c. $(-4x)(-3x + 2)$
 $= 12x^2 - 8x$


d. $(5x)(3x - 4)$
 $= 15x^2 - 20x$

4. Find the quotient.

a. $\frac{14n^2 - 2n}{2n}$
 $= 7n - 1$

b. $\frac{5x^2 - 3x}{-3}$
 $= \frac{5x^2}{-3} + x$

B 5. Which of the following quotients is represented in the tiles below?



A. $\frac{6x - 12}{6}$

B. $\frac{-6x + 12}{6}$

C. $\frac{-6x - 12}{6}$


Part 5: Surface Area of Composite Shapes

To determine the surface area of a 3D composite object, it is important to determine how the composite object is constructed.


- Using symmetry, determine similar faces.
- Determine the area of each face.
- Add these areas together.
- Consider how the composite shape is constructed and remove any overlapping areas.

Example 1

Consider the following 3D composite object.




This object is constructed of the following pieces:



Example 2

What is the surface area of the following shape?

**SA Piece A**

$$SA = 2(lh) + 2(lw) + 2(hw)$$

$$SA = 2(10 \times 2) + 2(3 \times 2) + 2(10 \times 3)$$

$$SA = 40 + 12 + 60$$

$$SA = 112 \text{ cm}^2$$

SA Piece B

$$SA = 2(lh) + 2(lw) + 2(hw)$$

$$SA = 2(15 \times 5) + 2(3 \times 5) + 2(15 \times 3)$$

$$SA = 150 + 30 + 90$$

$$SA = 270 \text{ cm}^2$$

Overlap on Piece A (Same Overlap on Piece B)

$$\text{Area} = lw$$

$$A = 3 \times 2$$

$$A = 6 \text{ cm}^2$$


$$\text{Total SA} = A + B - \text{Overlaps}$$

$$SA = 112 + 270 - 6 - 6$$


$$SA = 370 \text{ cm}^2$$

Practice Questions

1. Draw the objects that represent each composite.



2. Calculate the surface area of the following composite object.



SA Top Shape

$$\begin{aligned}
 SA &= 2(lh) + 2(lw) + 2(hw) \\
 SA &= 2(10 \times 5) + 2(10 \times 5) + 2(5 \times 5) \\
 SA &= 100 + 100 + 50 \\
 SA &= 250 \text{ cm}^2
 \end{aligned}$$

SA Base Shape $\times 2$

$$\begin{aligned}
 SA &= 2(lh) + 2(lw) + 2(hw) \\
 SA &= 2(2 \times 15) + 2(2 \times 2) + 2(15 \times 2) \\
 SA &= 60 + 8 + 60 \\
 SA &= 128 \text{ cm}^2 \times 2 = 256 \text{ cm}^2
 \end{aligned}$$

Overlap (on top shape and on column) $\times 2$ columns
Area = lw

$$\begin{aligned}
 A &= 2 \times 2 \\
 A &= 4 \text{ cm}^2 \times 2 = 8 \times 2 \text{ cm}^2 = 16 \text{ cm}^2
 \end{aligned}$$

Total SA = A + B - Overlaps

$$\begin{aligned}
 SA &= 250 + 256 - 16 \\
 SA &= 490 \text{ cm}^2
 \end{aligned}$$

